PLANT AND SOIL

Contents Volume 227, No. 1/2 2000

Water deficit and plant competition effects on growth and water-use efficiency of white clover (*Trifolium repens*, L.) and ryegrass (*Lolium perenne*, L.)
D.W. Lucero, P. Grieu and A. Guckert 1–15

Photosynthesis, respiration, and carbon allocation of two cool-season perennial grasses in response to surface soil drying
B. Huang and J. Fu 17–26

Remedying the iron-deficient maize plants by new synthetic macromolecular chelating agents
M. Ignatova, N. Manolova, I. Rashkov, V. Vassileva and G. Ignatov 27–34

Leachate from a Waste Disposal Centre reduces the initiation of arbuscular mycorrhiza, and spread of hyphae in soil
G.S. Pattinson, B.G. Sutton and P.A. McGee 35–45

A proposal for universal formulas for estimating leaf water status of herbaceous and woody plants based on spectral reflectance properties
G.-R. Yu, T. Miwa, K. Nakayama, N. Matsuoka and H. Kon 47–58

Stem tissue phosphorus as an index of the phosphorus status of *Banksia ericifolia* L. f.
S.E. Parks, A.M. Haigh, G.C. Cresswell 59–65

Allocation of carbon to shoots, roots, soil and rhizosphere respiration by barrel medic (*Medicago truncatula*) before and after defoliation
M.C. Crawford, P.R. Grace and J.M. Oades 67–75

Sap flow as an indicator of transpiration and the water status of young apricot trees
J.J. Alarcón, R. Domingo, S.R. Green, M.J. Sánchez-Blanco, P. Rodríguez and A. Torrecillas 77–85

Prediction of take-all disease risk in field soils using a rapid and quantitative DNA soil assay
Herdina and D.K. Roget 87–98

Effects of tillage intensity on nitrogen dynamics and productivity in legume-based grain systems
L.E. Drinkwater, R.R. Janke and L. Rossoni-Longnecker 99–113

Leaf endophyte *Neotyphodium coenophialum* modifies mineral uptake in tall fescue
D.P. Malinowski, G.A. Alloush and D.P. Belesky 115–126

Carbon dynamics and microbial activity in tallgrass prairie exposed to elevated CO_2 for 8 years
M.A. Williams, C.W. Rice and C.E. Owensby 127–137

The influence of temperature and nitrogen source on growth and nitrogen uptake of two polar-desert species, *Saxifraga caespitosa* and *Cerastium alpinum*
A. Volder, L.C. Bliss and H. Lambers 139–148

Competition in tree row agroforestry systems. 1. Distribution and dynamics of fine root length and biomass
S.J. Livesley, P.J. Gregory and R.J. Buresh 149–161

Biological control of root rot-root knot disease complex of tomato
I.A. Siddiqui, S.A. Qureshi, V. Sultana, S. Ehteshamul-Haque and A. Ghaffar 163–169

Below-ground nitrogen transfer between different grassland species: Direct quantification by ^{15}N leaf feeding compared with indirect dilution of soil ^{15}N
H. Høgh-Jensen and J.K. Schjoerring **171–183**

Comparative uptake of nitrate by intact seedlings of C_3 (barley) and C_4 (corn) plants: Effect of light and exogenously supplied sucrose
H.L. Sehtiya and S.S. Goyal **185–190**

Promotion of utilization of arbuscular mycorrhiza through reduced P fertilization 1. Bioassays in a growth chamber
H. Kahiluoto, E. Ketoja and M. Vestberg, **191–206**

Fate of polycyclic aromatic hydrocarbons (PAH) in the rhizosphere and mycorrhizosphere of ryegrass
Ph. Binet J.M. Portal and C. Leyval **207–213**

On the assessment of root and soil respiration for soils of different textures: interactions with soil moisture contents and soil CO_2 concentrations
T.J. Bouma and D.R. Bryla **215–221**

Effect of fluoride supply on fluoride concentrations in five pasture species: Levels required to reach phytotoxic or potentially zootoxic concentrations in plant tissue
D.P. Stevens, M.J. McLaughlin, P.J. Randall and G. Keerthisinghe **223–233**

Modelling zinc uptake by rice crop using a Barber-Cushman approach
T. Adhikari and R.K. Rattan **235–242**

Selenium accumulation by sequentially grown wheat and rice as influenced by gypsum application in a seleniferous soil
K.S. Dhillon and S.K. Dhillon **243–248**

Estimates of root system topology of five plant species grown at steady-state nutrition
A. Glimskär **249–256**

Availability of cadmium and zinc accumulated in the leaves of *Thlaspi caerulescens* incorporated into soil
K. Perronnet, C. Schwartz, E. Gérard and J.L. Morel **257–263**

The role of *Eriophorum vaginatum* in CH_4 flux from an ombrotrophic peatland
A.L. Greenup, M.A. Bradford, N.P. McNamara, P. Ineson and J.A. Lee **265–272**

Regulation of nitrogen partitioning in field-grown almond trees: Effects of fruit load and foliar nitrogen applications
F. Youssefi, S.A. Weinbaum and P.H. Brown **273–281**

Denitrification and N mineralization from hairy vetch (*Vicia villosa* Roth) and rye (*Secale cereale* L.) cover crop monocultures and bicultures
R.C. Rosecrance, G.W. McCarty, D.R. Shelton and J.R. Teasdale **283–290**

Soil mineral nitrogen availability was unaffected by elevated atmospheric pCO_2 in a four year old field experiment (Swiss FACE)
V. Gloser, M. Ježíková, A. Lüscher, M. Frehner, H. Blum, J. Nösberger and U.A. Hartwig **291–299**

Natural and induced cadmium-accumulation in poplar and willow: Implications for phytoremediation
B.H. Robinson, T.M. Mills, D. Petit, L.E. Fung, S.R. Green and B.E. Clothier **301–306**

Author index **307**